


# Chapter 12

## ADVANCED TRIGONOMETRY


### EXERCISE 12A

1 a




Data exhibits periodic behaviour.

b




Not enough information to say data is periodic. It may in fact be quadratic.

c




Not enough information to say data is periodic. It may in fact be quadratic.

d



Not enough information to say data is periodic.

2 a



c A curve can be fitted to the data as the distance travelled is continuous.


b The data is periodic.

- i The minimum value from the table is 0 and the maximum value is 64.  
So, the principal axis is  $y \approx \frac{0+64}{2}$ ,  
 $\therefore y \approx 32$ .
- ii The maximum value is  $\approx 64$  cm.
- iii The period is  $\approx 200$  cm.
- iv The amplitude is  $\approx 32$  cm.


3 a periodic    b periodic    c periodic    d not periodic    e periodic    f periodic

### EXERCISE 12B.1

- 1 a  $y = 3 \sin x$   
has amplitude 3 and period  $\frac{2\pi}{1} = 2\pi$   
When  $x = 0$ ,  $y = 0$ .



- b  $y = -3 \sin x$   
has amplitude  $|-3| = 3$   
and period  $\frac{2\pi}{1} = 2\pi$ .  
When  $x = 0$ ,  $y = 0$ .




It is the reflection of  $y = 3 \sin x$  in the  $x$ -axis.

**c**  $y = \frac{3}{2} \sin x$

has amplitude  $\frac{3}{2}$  and period  $\frac{2\pi}{1} = 2\pi$ .


When  $x = 0$ ,  $y = 0$ .



**d**  $y = -\frac{3}{2} \sin x$

has amplitude  $|\frac{-3}{2}| = \frac{3}{2}$

and period  $\frac{2\pi}{1} = 2\pi$ .




It is the reflection of  $y = \frac{3}{2} \sin x$  in the  $x$ -axis.

**2 a**  $y = \sin 3x$

has amplitude 1 and period  $\frac{2\pi}{3}$ .



When  $x = 0$ ,  $y = 0$ .



**c**  $y = \sin(-2x)$

has amplitude 1 and period  $\frac{2\pi}{|-2|} = \pi$ .

When  $x = 0$ ,  $y = 0$ .



It is the reflection of  $y = \sin 2x$  in the  $y$ -axis.

**3 a** period =  $\frac{2\pi}{|4|}$   
=  $\frac{\pi}{2}$

**b** period =  $\frac{2\pi}{|-4|}$   
=  $\frac{\pi}{2}$

**c** period =  $\frac{2\pi}{|\frac{1}{3}|}$   
=  $6\pi$

**d** period =  $\frac{2\pi}{0.6}$   
=  $\frac{20\pi}{6} = \frac{10\pi}{3}$

**4 a**  $\frac{2\pi}{B} = 5\pi$   
 $\therefore B = \frac{2}{5}$


**b**  $\frac{2\pi}{B} = \frac{2\pi}{3}$   
 $\therefore B = 3$

**c**  $\frac{2\pi}{B} = 12\pi$   
 $\therefore B = \frac{1}{6}$


**d**  $\frac{2\pi}{B} = 4$   
 $\therefore B = \frac{\pi}{2}$

**e**  $\frac{2\pi}{B} = 100$   
 $\therefore B = \frac{2\pi}{100} = \frac{\pi}{50}$


**5 a**





**b**



**c**




**EXERCISE 12B.2**


**3 a** period =  $\frac{2\pi}{|5|} = \frac{2\pi}{5}$       **b** period =  $\frac{2\pi}{|\frac{1}{4}|} = 8\pi$       **c** period =  $\frac{2\pi}{|-2|} = \pi$

**4 a**  $\frac{2\pi}{B} = 3\pi$       **b**  $\frac{2\pi}{B} = \frac{\pi}{10}$       **c**  $\frac{2\pi}{B} = 100\pi$       **d**  $\frac{2\pi}{B} = 50$


$\therefore B = \frac{2}{3}$        $\therefore B = 20$        $\therefore B = \frac{2}{100} = \frac{1}{50}$        $\therefore B = \frac{2\pi}{50} = \frac{\pi}{25}$

**5 a** A translation of  $\begin{pmatrix} 0 \\ -1 \end{pmatrix}$ , or vertically down 1 unit.

**b** A translation of  $\begin{pmatrix} \frac{\pi}{4} \\ 0 \end{pmatrix}$ , or horizontally  $\frac{\pi}{4}$  units right.



Month, $t$	1	2	3	4	5	6	7	8	9	10	11	12
Temp, $T$	0	-4	-10	-15	-16	-17	-18	-19	-17	-13	-6	-1



The period is  $\frac{2\pi}{B} = 12 \quad \therefore B = \frac{\pi}{6} \quad \{B > 0\}$

$$\text{Amplitude, } A \approx \frac{\text{max.} - \text{min.}}{2} \approx \frac{0 - (-19)}{2} \approx 9.5$$


$$D \approx \frac{\text{max.} + \text{min.}}{2} \approx \frac{0 + (-19)}{2} \approx -9.5$$

At min.,  $t = 8$  and at max.,  $t = 1 + 12 = 13 \quad \therefore C \approx \frac{8 + 13}{2} \approx 10.5$

$$\text{So, } T \approx 9.5 \sin \frac{\pi}{6}(t - 10.5) - 9.5 \quad \dots (1)$$

From technology,  $T \approx 13.1 \sin(0.345t + 2.37) - 5.43$

$$\therefore T \approx 13.1 \sin 0.345(t + 6.87) - 5.43 \quad \dots (2)$$



Neither model seems appropriate.

- 4 a For the model  $H = A \sin B(t - C) + D$

$$\text{period} = \frac{2\pi}{B} = 12.4 \text{ hours} \quad \therefore B = \frac{2\pi}{12.4} \approx 0.507$$

We let the principal axis be 0, so  $D = 0$


$\therefore$  the amplitude  $A = 7$ , so the min. is  $-7$ , and the max. is  $+7$

Let  $t = 0$  correspond to 'low tide'  $\therefore t = 6.2$  corresponds to 'high tide'

$$\therefore C = \frac{0 + 6.2}{2} = 3.1$$

$$\text{So, } H \approx 7 \sin 0.507(t - 3.1) + 0$$

$$\therefore H \approx 7 \sin 0.507(t - 3.1)$$



- 5 Let the model be  $H = A \sin B(t - C) + D$  metres

When  $t = 0$ ,  $H = 2$  and when  $t = 50$ ,  $H = 22$

$\uparrow$  min.  $\uparrow$  max.


$$\text{period} = \frac{2\pi}{B} = 100 \quad \therefore B = \frac{2\pi}{100} = \frac{\pi}{50}$$

$$A = 10 \quad \{\text{from the diagram}\} \quad D = \frac{\text{max.} + \text{min.}}{2} = \frac{22 + 2}{2} = 12$$

$$C = \frac{0 + 50}{2} = 25 \quad \{\text{values of } t \text{ at min. and max.}\} \quad \therefore H = 10 \sin \frac{\pi}{50}(t - 25) + 12$$


**EXERCISE 12D**

**1 a**  $y = \cos x + 2$




This is a vertical translation of  $y = \cos x$  through  $\begin{pmatrix} 0 \\ 2 \end{pmatrix}$ .

**b**  $y = \cos x - 1$




This is a vertical translation of  $y = \cos x$  through  $\begin{pmatrix} 0 \\ -1 \end{pmatrix}$ .

**c**  $y = \cos(x - \frac{\pi}{4})$




This is a horizontal translation of  $y = \cos x$  through  $\begin{pmatrix} \frac{\pi}{4} \\ 0 \end{pmatrix}$ .

**d**  $y = \cos(x + \frac{\pi}{6})$




This is a horizontal translation of  $y = \cos x$  through  $\begin{pmatrix} -\frac{\pi}{6} \\ 0 \end{pmatrix}$ .

**e**  $y = \frac{2}{3} \cos x$




This is a vertical stretch of  $y = \cos x$  with factor  $\frac{2}{3}$ .

**f**  $y = \frac{3}{2} \cos x$




This is a vertical stretch of  $y = \cos x$  with factor  $\frac{3}{2}$ .

**g**  $y = -\cos x$




This is a reflection of  $y = \cos x$  in the  $x$ -axis.

**h**  $y = \cos(x - \frac{\pi}{6}) + 1$




This is a translation of  $\begin{pmatrix} \frac{\pi}{6} \\ 1 \end{pmatrix}$ .

**i**  $y = \cos(x + \frac{\pi}{4}) - 1$




This is a translation of  $\begin{pmatrix} -\frac{\pi}{4} \\ -1 \end{pmatrix}$ .

**j**  $y = \cos(2x)$




This is a horizontal compression of factor 2.

**k**  $y = \cos(\frac{x}{2})$



This is a horizontal stretch of factor 2.

**l**  $y = 3 \cos(2x)$



This is a horizontal compression of factor 2 followed by a vertical stretch of factor 3.